



**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 12**

**SEPTEMBER 2023**

**AGRICULTURAL SCIENCES P1  
MARKING GUIDELINE**

**MARKS: 150**

---

This marking guideline consists of 11 pages.

---

**SECTION A****QUESTION 1**

1.1	1.1.1	B ✓✓		
	1.1.2	A ✓✓		
	1.1.3	D ✓✓		
	1.1.4	B ✓✓		
	1.1.5	C ✓✓		
	1.1.6	B ✓✓		
	1.1.7	C ✓✓		
	1.1.8	A ✓✓		
	1.1.9	C ✓✓		
	1.1.10	D ✓✓	(10 x 2)	(20)
1.2	1.2.1	Both A and B ✓✓		
	1.2.2	A only ✓✓		
	1.2.3	B only ✓✓		
	1.2.4	None ✓✓		
	1.2.5	A only ✓✓	(5 x 2)	(10)
1.3	1.3.1	Biological value ✓✓		
	1.3.2	Subsistence ✓✓		
	1.3.3	Dystocia ✓✓		
	1.3.4	Lymphatic system ✓✓		
	1.3.5	Lack of libido ✓✓	(5 x 2)	(10)
1.4	1.4.1	Fodder flow ✓		
	1.4.2	Chronic ✓		
	1.4.3	Scrotum ✓		
	1.4.4	Placenta ✓		
	1.4.5	Implantation ✓	(5 x 1)	(5)
			<b>TOTAL SECTION A:</b>	<b>45</b>


**SECTION B****QUESTION 2: ANIMAL NUTRITION****2.1 Alimentary canal of farm animals**

- 2.1.1 **Name of the part**  
Small intestines ✓ (1)
- 2.1.2 **TWO visible adaptation features**
- Presence of micro-villi ✓
  - Presence of blood capillaries ✓
  - Presence of lymph vessels ✓
- (Any 2 x 1) (2)
- 2.1.3 **Indication of nutrients absorbed in:**
- (a) **Lymph** – Digested fats ✓ (1)
- (b) **Blood capillaries** – Digested carbohydrates ✓ (1)
- 2.1.4 **Explanation of how folds assist in absorption**  
Folds increase the surface area ✓ for absorption ✓ (2)

**2.2 Feed components**

- 2.2.1 **Identification of**
- (a) **Energy – rich concentrate** – Maize meal ✓ (1)
- (b) **Protein – rich roughage** – Lucerne hay ✓ (1)
- 2.2.2 **Type of an animal**  
Ruminant ✓ (1)
- 2.2.3 **Reason**
- The ruminant animal has micro-organisms ✓ which are able to digest a ration containing roughage and urea ✓
  - The ruminant animal is able to regurgitate ✓ the feed for re-chewing of roughage ✓
  - The ruminant animal has four compartments ✓ in its stomach adapted to digest roughages ✓ (Any 1) (2)
- 2.2.4 **Component of the ration that can improve palatability and digestibility of oat hay**  
Molasses ✓ (1)

## 2.2.5 Bar graph

**CRITERIA/RUBRIC/MARKING GUIDELINE**

- Correct heading ✓
- Bar graph ✓
- x-axis: Correctly calibrated and labelled (Feed components) ✓
- y-axis : Correctly calibrated and labelled (Quantities) ✓
- Correct unit (kg) ✓
- Accuracy (80% + correctly plotted) ✓ (6)

## 2.3 Sow and its litter housed in a farrowing pen with a cement floor

2.3.1 **Mineral element deficient in sow**  
Iron/Fe ✓ (1)

2.3.2 **ONE deficiency symptom of iron**

- Anaemia ✓
- Paleness of the mucous membrane ✓
- Listlessness/laziness/fatigue ✓ (Any 1 x 1) (1)

2.3.3 **Method of supplementing iron**

- Injection ✓
- Soil sods placed in pig's concrete pen ✓
- Feeding with green forage ✓ (Any 1 x 1) (1)

## 2.4 Pearson square

### 2.4.1 Ratio representing sunflower oilcake meal

8 ✓ (1)

### 2.4.2 Reason

A feed high in protein ✓ constitutes a small part of the ratio in the mixture. ✓ (2)

### 2.4.3 Calculation of the percentage of a carbohydrate-rich feed in the mixture.

$8 + 20 = 28$  ✓  
 $= \frac{20}{28} \times 100$  ✓  
 $= 71,43\%$  ✓ (3)

## 2.5 Energy values of a feed

### 2.5.1 Identify the energy loss in B

Energy lost as body heat ✓ (1)

### 2.5.2 Justification of the importance of net energy

- Needed for production/growth/reproduction ✓
- Needed for maintenance ✓ (2)

### 2.5.3 Calculation of metabolic energy

Metabolic energy/ME =  
 = Gross energy – energy loss in faeces – energy loss in urine and fermentation gases  
 = 24J – 9J – 5J ✓  
 = 10J ✓

OR

= Digestible energy – energy loss in urine and fermentation gases  
 = 15J – 5J ✓  
 = 10J ✓ (2)

### 2.5.4 TWO aims of calculating energy value of the feed

- To determine the animal's diet ✓
- To determine feeding standards ✓
- To determine ration formulation ✓

(Any 2 x 1) (2)

**[35]**

**QUESTION 3: ANIMAL PRODUCTION, PROTECTION AND CONTROL****3.1 Handling facilities**

- 3.1.1 **Identification of the facility**  
Loading ramp ✓ (1)
- 3.1.2 **Purpose of facility labelled A/crush**
- To restrain farm animals/ ✓
  - To guide farm animals to the vehicle for transportation ✓ (1)
- 3.1.3 **Design feature of a crush to ensure safety**
- Has high solid/strong/solid sides to prevent animals from seeing out ✓
  - It has curves that are not sharp ✓
  - There is nothing that can harm/hurt animals ✓
  - Angles are not too steep ✓ (Any 2 x 1) (2)
- 3.1.4 **TWO reasons for handling farm animals using crush**
- For animal health programmes ✓
  - Normal management programme/dehorning/castration/markings/docking ✓
  - Treatment of parasites ✓
  - Determination of animal's age ✓
  - Generation of data ✓
  - Transportation of animals ✓ (Any 2 x 1) (2)

**3.2. Animal handling**

- 3.2.1 **Indication of the letter**
- (a) A ✓ (1)
  - (b) D ✓ (1)
  - (c) C ✓ (1)
- 3.2.2 **Behaviour when approached at blind spot**
- It will kick
  - It will be restless/uncomfortable ✓ (1)
- 3.2.3 **TWO common behaviours displayed by cattle under stress**
- Pinned or raised ears ✓
  - Rapid tail movement ✓
  - Raised hair on the back of the neck ✓
  - Pawing ✓
  - Snorting ✓
  - Feigned charging movements ✓ (Any 2 x 1) (2)

### 3.3 Shelter and housing farm animals

#### 3.3.1 Purpose of using the structures

- (a) **Holding pen** – For keeping animals temporarily prior to handling ✓
- (b) **Farrowing pen** – Keeping sows and piglets ✓
- (c) **Holding shed** – Keeping animals for a long period of time to protect them against temperature changes ✓ (3)

#### 3.3.2 THREE reasons for shelter/housing in animal production

- To protect animals against extreme temperature changes ✓
- To protect animals from predators/thieves ✓
- For easy handling ✓ (3)

### 3.4 Animal diseases

#### 3.4.1 Labelling

- A** – Rabies ✓
- B** – Bacteria ✓
- C** – Swollen udder ✓  
– Milk is thick, flaky with clots ✓ (Any 1)
- D** – Coccidiosis ✓
- E** – Hygiene ✓ (5)

#### 3.4.2 Identification of the role of

- (a) The farmer – Good hygienic principles ✓ (1)
- (b) The state – Provision of immunisation/vaccination ✓ (1)

### 3.5 Methods of administering medicine to animals

#### 3.5.1 Identification of methods to apply medicine

- A** – Topical ✓
- B** – Vaginal insertion ✓
- D** – Plunge dipping/Dipping ✓ (3)

#### 3.5.2 Letter representing the method used to treat parasites

- (a) **Roundworm** – C ✓ (1)
- (b) **Blue ticks** – D ✓ (1)

**3.5.3 TWO ways of using medication sustainably**

- Medicine is safe to use for the specific animal ✓
- Check the expiry date ✓
- Ensure proper storage ✓
- Administer correct dose ✓
- Administer according to the instructions ✓
- Administer medicine for the correct period to ensure its effectiveness ✓
- Allow for proper withdrawal period before it is consumed ✓
- Medicine be kept away from children ✓ (Any 2 x 1) (2)

**3.6 Poisonous plants****3.6.1 ONE poisonous plant found in pastures**

- Thorn apple ✓
- Poisonous bulb ✓
- Lantana ✓
- Devil's thorn ✓
- Lupines ✓
- Buffalo grass ✓
- Poisonous leaf ✓ (Any 1 x 1) (1)

**3.6.2 TWO measures to control plant poison in pastures**

- Remove animals from an infested camp ✓
  - Remove poisonous plants from the pastures ✓
  - Feed animal well as they will be less likely to eat poisonous plants ✓
  - Avoid overgrazing/overstocking ✓
  - Practise rotational grazing ✓
  - Provide animals with feed and water when transported by rail/when introducing them to a new place with unfamiliar plants ✓
  - Control poisonous plants by applying chemicals in infested pastures ✓ (Any 2) (2)
- [35]**



**QUESTION 4: ANIMAL REPRODUCTION****4.1 Reproductive system****4.1.1 Identification of the letter**

- (a) B ✓ (1)  
(b) D ✓ (1)  
(c) E ✓ (1)

**4.1.2 TWO congenital defects in part B/testis leading to loss of fertility**

- Hypoplasia ✓
- Cryptorchidism ✓
- Sperm defect ✓ (Any 2 x 1) (2)

**4.1.3 Role played by part labelled C/penis in reproduction**

It deposits semen into the vagina during mating ✓ (1)

**4.2 Hormones controlling oestrus cycle****4.2.1 Duration of the oestrus**

28 days ✓ (1)

**4.2.2 Name of the hormones**

**A** – Oestrogen ✓  
**B** – Progesterone ✓ (2)

**4.2.3 Indication of what is happening during the follicular phase**

- (a) Stage of oestrus cycle – Pro-oestrus ✓  
(b) Hormone responsible – FSH ✓ (2)

**4.2.4 TWO functions of hormone B/progesterone if the cow can be pregnant.**

- Delays secretion of FSH ✓
- Prevents the cow from coming to heat ✓
- Prepares the uterus to receive the fertilised egg ✓
- Maintains proper uterine environment to maintain pregnancy ✓
- Stimulating uterine milk secretions ✓ (Any 2 x 1) (2)

### 4.3 Artificial insemination

#### 4.3.1 Purpose of using the pistolette

To deposit semen during artificial insemination ✓ (1)

#### 4.3.2 TWO basic requirements for storage

- Semen to be stored at 5°C if stored for a short period ✓
- Semen be kept frozen in liquid nitrogen at -196°C if stored for a longer time ✓
- Must be stored in polyvinyl straws ✓
- The ends of straws are sealed to prevent liquid nitrogen from entering ✓
- Straws should be labelled for identification ✓ (Any 2 x 1) (2)

#### 4.3.3 Identification of the letter

- (a) A ✓ (1)  
(b) E ✓ (1)

#### 4.3.4 ONE disadvantage of using the equipment for the farmer

It is expensive ✓ (1)

#### 4.3.5 TWO advantages of artificial insemination

- Decreases the occurrence of sexually transmitted diseases ✓
- More female animals can be fertilised by superior male animals ✓
- It is a quick and economical way to improve the herd ✓
- Semen from males in other countries can be used ✓
- Semen of superior bulls can be used even after death ✓
- It improves the commercial value of the herd ✓
- Semen of multiple sires can be used without keeping and maintaining expensive bulls ✓ (Any 2 x 1) (2)

### 4.4 Embryo harvesting/flushing

#### 4.4.1 Identification of the procedure

Embryo harvesting/flushing ✓ (1)

#### 4.4.2 Type of a cow where the procedure is performed

Donor/superior cow ✓ (1)

#### 4.4.3 Reason

It possesses genetically superior desirable characteristics ✓ (1)

**4.4.4 ONE aim of embryo transplant/ET**

- To prevent extinction of valuable animals / increase the number of endangered species ✓
- To improve disease resistance by using embryos of superior animals that are resistant to certain diseases ✓
- To improve the growth rate and production yields ✓ (Any 1 x 1) (1)

**4.4.5 TWO disadvantages of the technique for the farmer**

- More expensive/labour intensive ✓
  - Needs considerable skill and experience ✓
  - Synchronisation of the recipient and donor can be difficult ✓
  - Recipient cow may not become pregnant/abortion may occur ✓
  - Recipients may not have a strong enough heat cycle to accept the insemination ✓
  - There is a danger that recipients could abort the embryos ✓
  - Time consuming
- (Any 2 x 1) (2)

**4.5 Flow of milk****4.5.1 Rearrangement of the steps with which the milk will flow**

- Alveolus ✓
  - Milk ducts ✓
  - Gland cistern ✓
  - Teat cistern ✓
  - Teat canal ✓
- (5)

**4.5.2 Milk let down process****(a) TWO stimuli that initiate the process**

- Washing of the udder ✓
- Massage of the udder ✓
- Milking action ✓
- Appearance and sound of the milker ✓
- Seeing the calf ✓ (Any 2 x 1) (2)

**(b) Hormone involved**

- Oxytocin ✓ (1)

**[35]****TOTAL SECTION B: 105****GRAND TOTAL: 150**