



Province of the  
**EASTERN CAPE**  
EDUCATION



**NATIONAL  
SENIOR CERTIFICATE/  
NASIONALE  
SENIORSERTIFIKAAT**

**GRADE/GRAAD 12**

**SEPTEMBER 2023**

**PHYSICAL SCIENCES P2/ FISIESE  
WETENSKAPPE V2  
MARKING GUIDELINE/NASIENRIGLYN**

**MARKS: 150**

---

This marking guideline consists of 19 pages./  
*Hierdie nasienriglyn bestaan uit 19 bladsye.*

---

**QUESTION 1/VRAAG 1**

- |      |      |     |
|------|------|-----|
| 1.1  | D ✓✓ | (2) |
| 1.2  | B ✓✓ | (2) |
| 1.3  | A✓✓  | (2) |
| 1.4  | D ✓✓ | (2) |
| 1.5  | B ✓✓ | (2) |
| 1.6  | C ✓✓ | (2) |
| 1.7  | D ✓✓ | (2) |
| 1.8  | A ✓✓ | (2) |
| 1.9  | D ✓✓ | (2) |
| 1.10 | B ✓✓ | (2) |
- [20]**

**QUESTION 2/VRAAG 2**

- 2.1.1 D ✓ (1)  
 2.1.2 F ✓ (1)  
 2.1.3 B ✓ (1)  
 2.2 UNSATURATED ✓


It is an organic compound that contains double/triple/multiple bonds. ✓

ONVERSADIG

*Dit is 'n organiese verbinding wat dubbel/drievoudige/meervoudige bindings bevat.*

(2)

2.3.1



**Marking criteria/Nasienkriteria**

- Functional group correct ✓  
*Funksionele groep korrek*
- Whole structure correct ✓  
*Hele struktuur korrek*

(2)

2.3.2 hept-3-yne ✓✓ / 3-heptyne

hept-3-yn / 3-heptyn

**Marking criteria/ Nasienkriteria**

- Parent name and suffix correct (heptyne)✓  
*Stam naam en agtervoegsel korrek (heptyn)*
- Everything correct e.g. hyphens and numbering ✓  
*Alles korrek bv koppeltekens en nommering*

(2)

2.4 Tertiary. ✓

The carbon that is bonded to the hydroxyl group (-OH) is bonded to three other carbons. ✓

**OR**

The carbon of the functional group is bonded to three other carbons.

*Tertiêre*

*Die koolstof wat verbind is aan die hidroksielgroep (-OH) is verbind aan drie ander koolstowwe.*

**OF**


*Die koolstof van die funksionele groep is aan drie ander koolstowwe verbind.* (2)

2.5 Butan-2-ol / 2-butanol ✓✓ **NOTE/LET WEL:** Butan-1-ol / 1-butanol (1/2) (2)

2.6.1 Compounds with the same molecular formula ✓ but different functional groups ✓/belong to different homologous series.

*Verbindings met dieselfde molekulêre formule, maar verskillende funksonale groepe / behoort aan verskillende homoloë reeks.* (2)

2.6.2 CH<sub>3</sub>COCH<sub>3</sub> ✓✓      OR/OF



(2)  
[17]

### QUESTION 3/VRAAG 3

#### 3.1 **Marking criteria/ Nasienriglyne**

If any of the underlined key words/phrases in the **correct context** are omitted: -1 mark per word/phrase.

*Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: -1 punt per woord/frase*

The temperature at which the vapour pressure of a liquid equals the atmospheric pressure ✓✓

*Die temperatuur waarby die dampdruk van 'n vloeistof gelyk is aan die atmosferiese druk.* (2)

3.2.1 Molecular size ✓ / Surface area / Chain length / London forces  
*Molekulêre grootte / Oppervlakte / kettinglengte / Londonkragte* (1)

3.2.2 Functional group ✓/ Homologous series  
*Funksionele groep / Homoloë reeks* (1)

3.3 London forces / induced dipole forces / dispersion forces ✓  
*Londonkragte / geïnduseerde dipoolkragte / verspreidingskragte* (1)

3.4 129 °C ✓ (1)

3.5

**Marking criteria/Nasienkriteria**

- Compare the molecular size of 2-methylbutan-1-ol to butan-1-ol ✓
- Relate the molecular size to London forces/induced dipole forces/dispersion forces. ✓
- Compare the chain length of 2-methylbutan-1-ol to pentan-1-ol ✓
- Relate the chain length to London forces/induced dipole forces/dispersion forces ✓
- *Vergelyk die molekulêre grootte van 2-metielbutan-1-ol aan butan-1-ol*
- *Verwys die molekulêre grootte na die Londonkragte/ geïnduseerde dipoolkragte / verspreidingkragte*
- *Vergelyk die kettinglengte van 2-metielbutan-1-ol met pentan-1-ol*
- *Verwys die kettinglengte na die Londonkragte / geïnduseerde dipoolkragte / verspreidingkragte*

- 2-methylbutan-1-ol has a larger molar mass/molecular size than butan-1-ol ✓
- London forces/induced dipole forces/dispersion forces of 2-methylbutan-1-ol is stronger than that of butan-1-ol ✓
- The boiling point will be higher than that of butan-1-ol
- 2-methyl butan-1-ol has a shorter chain length than pentan-1-ol ✓
- London forces of 2-methyl butan-1-ol is weaker than that of pentan-1-ol ✓
- The boiling point will be lower than that of pentan-1-ol
- *2-metielbutan-1-ol het 'n groter molekulêre massa/molekulêre grootte as butan-1-ol*
- *Londonkragte/geïnduseerde dipoolkragte/verspreidings van 2-metielbutan-1-ol is sterker as dié van butan-1-ol*
- *Die kookpunt is hoër as dié van butan-1-ol*
- *2-metielbutan-1-ol het 'n korter kettinglengte as pentan-1-ol*
- *Londonkragte van 2-metielbutan-1-ol is swakker as dié van pentan-1-ol*
- *Die kookpunt sal laer wees as dié van pentan-1-ol*

(4)

## OR /OF

- Butan-1-ol has a smaller molar mass/molecular size than 2-methylbutan-1-ol
- London forces/induced dipole forces/dispersion forces of butan-1-ol is weaker than that of 2-methylbutan-1-ol
- The boiling point will be higher than that of butan-1-ol
- Pentan-1-ol has a larger chainlength than 2-methyl butan-1-ol
- London forces/induced dipole forces/dispersion forces of pentan-1-ol is stronger than that of 2-methyl butan-1-ol
- The boiling point will be lower than that of pentan-1-ol
- *Butan-1-ol het 'n kleiner molekulêre massa/molekulêre grootte as 2-metielbutan-1-ol*
- *Londonkragte/geïnduseerde dipoolkragte/verspreidings van butan-1-ol is swakker as dié van 2-metielbutan-1-ol*
- *Die kookpunt is hoër as dié van butan-1-ol*
- *Pantan-1-ol het 'n langer kettinglengte as 2-metielbutan-1-ol*
- *Londonkragte van pentan-1-ol is sterker as dié van 2-metielbutan-1-ol*
- *Die kookpunt sal laer wees as dié van pentan-1-ol*

3.6.1

**Marking criteria/Nasienkriteria**If any of the underlined key words/phrases in the **correct context** are omitted:

- 1 mark per word/phrase.

*Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: - 1 punt per woord/frase*

The pressure exerted by a vapour at equilibrium with its liquid in a closed system. ✓✓

*Die druk uitgeoefen deur 'n damp in ewewig met sy vloeistof in 'n geslote sisteem.* (2)

3.6.2

**Q ✓**

(1)

3.6.3

**Marking criteria/Nasienkriteria**

- Propan-1-ol has hydrogen bonds ✓(and London forces/induced dipole forces/dispersion forces)
- Propanal has dipole-dipole forces ✓( and London/induced dipole forces/dispersion forces)
- Compare the strength of the hydrogen bonds to dipole-dipole forces ✓
- Relate strength of intermolecular forces to vapour pressure ✓
- *Propan-1-ol het waterstofbindings (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Propanal het dipool-dipoolkragte (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Vergelyk die sterkte van die waterstofbindings met die dipool-dipoolkragte*
- *Verwys die sterkte van die intermolekulêrekragte met die dampdruk*

- Propan-1-ol has hydrogen bonds ✓ (and London forces/induced dipole forces/dispersion forces)
- Propanal has dipole-dipole forces (and London forces/induced dipole forces/dispersion forces) ✓
- Hydrogen bonds are stronger than the dipole-dipole forces ✓
- Stronger intermolecular forces result in lower vapour pressure ✓
  
- *Propan-1-ol het waterstofbindings (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Propanal het dipool-dipoolkragte (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Waterstofbindings is sterker as dipool-dipoolkragte*
- *Sterker intermolekulêrekragte het laer dampdruk*

**OR / OF**

- Propan-1-ol has Hydrogen bonds ✓ (and London forces/induced dipole forces/dispersion forces)
- Propanal has dipole-dipole forces ✓ ( and London/induced dipole forces/dispersion forces)
- Dipole-dipole forces are weaker than the hydrogen bonds✓
- Weaker intermolecular forces result in higher vapour pressure ✓
  
- *Propan-1-ol het waterstofbindings (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Propanal het dipool-dipoolkragte (en Londonkragte/geïnduseerde dipoolkragte/verspreidingskragte)*
- *Dipool-dipoolkragte is swakker as waterstofbindings*
- *Swakker intermolekulêrekragte het hoër dampdruk*

(4)

[17]


**QUESTION 4/VRAAG 4**

4.1.1 Hydrolysis / *Hidrolise* ✓ (1)

4.1.2 Pentan-2-ol / 2-pentanol ✓✓ (2)

4.1.3 Concentrated strong base ✓ OR concentrated NaOH OR concentrated KOH OR concentrated LiOH  
*Gekonsentreerde sterk basis OF gekonsentreerde NaOH OF gekonsentreerde KOH OF gekonsentreerde LiOH* (1)

4.1.4

**Marking criteria/Nasienkriteria**

- Functional group correct ✓  
*Funksionele groep korrek*
- Whole structure correct ✓  
*Hele struktuur korrek*

(2)


4.1.5 Sulphuric acid / *Swawelsuur* ✓ (1)

4.1.6 Dehydration / *Dehidrasie* / *Dehidratering* ✓ (1)

4.2.1 Esterification / Condensation / *Esterifikasie* / *Veresterung* ✓ (1)

4.2.2 Alcohols are flammable✓ / prevent fire  
*Alkohole is vlamaar / om 'n vuur te voorkom* (1)

4.2.3

**Marking criteria/Nasienkriteria**

- Functional group correct ✓  
*Funksionele groep korrek*
- Whole structure correct ✓  
*Hele struktuur korrek*

ethyl ✓propanoate ✓ / *etiel-propanoaat* (4)

[14]

**QUESTION 5/VRAAG 5**

5.1

**Marking criteria/ Nasienriglyne**

If any of the underlined key words/phrases in the **correct context** are omitted: - 1 mark per word/phrase.

*Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: - 1 punt per woord/frase*

**ANY ONE**

Change in concentration ✓ of reactant or product per (unit) time.✓

Change in amount/number of moles/volume/mass of products or reactants per (unit) time.

Change in amount/number of moles/volume/mass of products formed or reactants used reactants per (unit) time.

**ENIGE EEN**

Verandering in konsentrasie van reaktanse of produkte per (eenheid) tyd.

Verandering in hoeveelheid/getal mol/volume/massa van reaktanse of produkte per (eenheid) tyd.

Verandering in hoeveelheid/getal mol/volume/massa van produkte gevorm/reaktanse gebruik per (eenheid) tyd.

**OR/OF**

The rate of change in concentration / amount of moles / number of moles / volume / mass. **(2 or 0)**

*Die tempo van verandering in konsentrasie/hoeveelheid mol/getal mol/volume/massa ✓✓ **(2 of 0)***

(2)

5.2

Stopwatch ✓/Timer/ Measuring cylinder

*Stophorlosie / Tydhouer / Meetsilinder*

(1)

5.3

Reaction is exothermic / reaction releases heat ✓

*Reaksie is eksotermies / reaksie gee hitte af*

(1)

5.4

SHORTER / KORTER ✓

(1)

## 5.5 For Exp. 2

- Higher concentration results in more particles colliding with correct orientation ✓
- The number of effective collision per unit time increases / frequency of the effective collisions increases ✓

## Vir Eksp. 2

- Hoër konsentrasie beteken meer deeltjies bots met die korrekte orientasie
- Die aantal effektiewe botsings per eenheid tyd neem toe/ frekwensie van die effektiewe botsings neem toe.

## OR / OF

## For Exp. 1

- Low concentration results in fewer particles colliding with correct orientation ✓
- The number of effective collision per unit time decreases / frequency of effective collisions decreases ✓

## Vir Eksp. 1

- Lae konsentrasie beteken minder deeltjies bots met die korrekte orientasie
- Die aantal effektiewe botsings per eenheid tyd neem af / frekwensie van die effektiewe botsings neem af.

(2)

5.6.1 Rate/ Tempo =  $\frac{\Delta c}{\Delta t}$

$$= \frac{250-0}{5,28-0} \checkmark \quad \text{Accept / Aanvaar} \quad \frac{250}{5,28}$$

$$= 0,95 \checkmark (\text{cm}^3 \cdot \text{min}^{-1}) \quad (3)$$

5.6.2 Marking criteria / Nasienkriteria

- $V_{\text{CO}_2}$  remaining / oorbly =  $150 \text{ cm}^3$
- Substitution into / Vervanging in  $n = V/V_m$
- Subst. into / Vervanging in  $n = m/M$
- Final answer / Finale antwoord

$$V(\text{CO}_2) \text{ remaining / oorbly} = 250 - 100 = 150 \text{ cm}^3 \checkmark$$

$$n = V/V_m = 150/25\ 000 \checkmark$$

$$= 6 \times 10^{-3} \text{ mol}$$

$$n(\text{CO}_2) = m/M$$

$$6 \times 10^{-3} = m / 44 \checkmark$$

$$m = 0,264 \text{ g } \checkmark / 0,26 \text{ g}$$

(4)

5.7.1 P ✓ (1)

5.7.2  $C_2H_4O_2$  is a weaker acid ✓ (than  $HCl$ ) which will result in a LOWER reaction rate ✓ (for Expt. 3) / Lower gradient/Longer reaction time

$C_2H_4O_2$  is 'n swakker suur (as  $HCl$ ) wat sal lei na 'n LAER reaksietempo (vir Eksp. 3) / laer gradiënt / langer reaksietyd

**OR / OF**

$HCl$  is stronger acid ✓ (than  $C_2H_4O_2$ ) which will result in a HIGHER reaction rate ✓ for Expt. 1/ Higher gradient/Shorter reaction time

$HCl$  is 'n sterker suur (as  $C_2H_4O_2$ ) wat sal lei na 'n HOËR reaksietempo (vir Eksp. 1) / hoër gradiënt / korter reaksietyd (2)

5.7.3 EQUAL TO ✓  
Final amount of  $CO_2$  is the same ✓ (in both experiments)

**GELYK AAN**

*Finale hoeveelheid van  $CO_2$  is gelyk (in beide eksperimente)*

(2)

[19]

**QUESTION 6 / VRAAG 6**

- 6.1.1 Reversible reaction ✓ /Products can be converted back to reactants  
*Omkeerbare reaksie / Produkte kan na reaktanse omgeskakel word*

**NOTE: Do not accept “Reaction is at equilibrium”**

**LET WEL: Moenie “Reaksie is by ewewig” aanvaar nie.**

(1)

- 6.1.2 (Chemical) Equilibrium / (*Chemiese*) Ewewig ✓

(1)

- 6.1.3 HEATED / VERHIT ✓

(1)

- 6.1.4 ENDOTHERMIC / ENDOTERMIES ✓


(1)

- 6.1.5 • Increase in temperature favoured the forward reaction ✓ / The rate of the forward reaction increased more than the rate of the reverse reaction.  
 • Increase in temperature favours the endothermic reaction. ✓

- *Toename in temperatuur bevoordeel die voortwaartse reaksie / Die tempo van die voortwaartse reaksie is hoër as die terugwaartse reaksie*
- *'n Toename in temperatuur bevoordeel die endotermiese reaksie*

(2)

- 6.1.6

**Marking criteria/ Nasienkriteria**

- Both rates increase; new equilibrium at a higher rate when P is increased ✓  
*Beide tempos neem toe; nuwe ewewig is by 'n hoër tempo wanneer P verhoog*
- The increase in reverse reaction is HIGHER than for the forward reaction and eventually there is horizontal section for both ✓  
*Die toename in die terugwaartse reaksie is HOËR as die voortwaartse reaksie en daar is horizontale gedeelte by die einde vir beide.*

**NOTE: Do not penalise if axes are not labelled.**

**LET WEL: Moenie penaliseer as asse nie benoem is nie.**

(2)

## 6.2 MOLE CALCULATIONS / MOL BEREKENINGE

- a. Correct  $K_c$  expression (formula in square brackets) ✓
- b. Substitution of equilibrium concentration of  $\text{CO}_2$  into  $K_c$  expression ✓
- c. Determining the  $n_{\text{equilibrium}}$  CO ✓
- d. Determining the  $n \text{ CO}_{\text{reacted}}$
- e. Correct mol ratio for  $\text{CO:C}$  ✓
- f. Determining the initial mol of C✓
- g. Correct substitution into percentage formula ✓
- h. Final answer ✓
  
- a. Korrekte  $K_c$ -uitdrukking (Formule formule tussen vierkantbakies)
- b. Vervanging van ewewigkonsentrasie van  $\text{CO}_2$  in  $K_c$  – uitdrukking
- c. Bepaal die  $n_{\text{ewewig}}$  CO
- d. Bepaal  $n \text{ CO}_{\text{reageer}}$
- e. Korrekte molverhouding vir  $\text{CO} : \text{C}$
- f. Bepaal die aanvanklike mol van C
- g. Korrekte vervanging in persentasie formule
- h. Finale antwoord

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \checkmark \text{ (a)}$$

$$0,05 = \frac{[\text{CO}]^2}{0,05} \checkmark \text{ (b)}$$

$$[\text{CO}] = 0,05 \text{ mol} \cdot \text{dm}^{-3}$$

- No  $K_c$  expression, correct substitution / Geen  $K_c$ -uitdrukking, korrekte, korrekte substitusie. Max. / Maks. 7/8
- Wrong  $K_c$  expression / Verkeerde  $K_c$ -uitdrukking. Max. Maks. 5/8

	C (s)	$\text{CO}_2$ (g)	2 CO (g)
Initial mol / Aanvangs-hoeveelheid (mol)	0,45 ✓ (f)		0
Change/ Verandering in mol	0,05 ✓ (e)		0,1 ✓ (d)
Equilibrium/ Ewewig mol	0,40		0,1 ✓ (c)
Concentration/ Konsentrasie (mol · dm <sup>-3</sup> )	-		0,05

$$\% \text{ C reacted} = \Delta n/n_{\text{initial}} \times 100$$

$$= 0,05/0,45 \times 100 \checkmark \text{ (g)}$$

$$= 11,11\% \checkmark \text{ (h)}$$

(8)  
[16]

**QUESTION 7/VRAAG 7**

- 7.1.1 An acid is a proton ( $H^+$ -ion) donor ✓✓  
*'n Suur is 'n proton ( $H^+$ -ion) skenker* (2)
- 7.1.2  $HC_2O_4^-$  ✓ (1)
- 7.1.3 Weak acid / *Swaksuur* ✓ (1)
- 7.1.4 The acid did not fully/completely ionise ✓✓ (in water). **OR** ionised incompletely.  
*Die suur ioniseer nie volledig (in water) nie **OF** Suur ioniseer onvolledig* (2)

**7.2.1 Marking criteria/Nasienkriteria**

- $n = cV$
- Subst. into / *Vervanging* in  $n = cV$
- Final answer / *Finale antwoord*

$$\begin{aligned} n (\text{CH}_3\text{COOH}) &= cV \checkmark \\ &= 0,1 \times 25 \times 10^{-3} \checkmark \\ &= 2,5 \times 10^{-3} \text{ mol} \checkmark \end{aligned}$$

(3)

<b>7.2.2</b>	<b><u>Marking criteria/Nasienkriteria</u></b>	<b><u>Marking criteria/Nasienkriteria</u></b>
	<ul style="list-style-type: none"> <li>• Formula <math>pH = -\log [H_3O^+]</math> ✓</li> <li>• pH value substituted into formula ✓</li> <li>• Substitution in <math>K_w</math> formula ✓</li> <li>• Final answer ✓</li>   <li>• <i>Formule pH = - log [H<sub>3</sub>O<sup>+</sup>]</i></li> <li>• <i>pH waarde vervang in formule</i></li> <li>• <i>Vervang in K<sub>w</sub> formule</i></li> <li>• <i>Finale antwoord</i></li> </ul>	<ul style="list-style-type: none"> <li>• Formula <math>pOH + pH = 14</math> ✓</li> <li>• pH value substituted into formula ✓</li> <li>• Substitution in pOH formula ✓</li> <li>• Final answer ✓</li>   <li>• <i>Formule pOH + pH = 14</i></li> <li>• <i>pH waarde vervang in formule</i></li> <li>• <i>Vervang van pOH waarde in formule</i></li> <li>• <i>Finale antwoord</i></li> </ul>

<u>OPTION 1 / OPSIE 1</u>	<u>OPTION 2 / OPSIE 2</u>
$\text{pH} = -\log [\text{H}_3\text{O}^+] \checkmark$	$\text{pOH} + \text{pH} = 14 \checkmark$
$12 \checkmark = -\log [\text{H}_3\text{O}^+]$	$\text{pOH} + 12 \checkmark = 14$
$[\text{H}_3\text{O}^+] = 1 \times 10^{-12} \text{ mol}\cdot\text{dm}^{-3}$	$\text{pOH} = 2$
$K_w = [\text{OH}^-][\text{H}_3\text{O}^+] = 1 \times 10^{-14}$	$\text{pOH} = -\log [\text{OH}^-]$
$[\text{OH}^-][\text{H}_3\text{O}^+] = 1 \times 10^{-14}$	$2 = -\log [\text{OH}^-] \checkmark$
$[\text{OH}^-](1 \times 10^{-12}) = 1 \times 10^{-14} \checkmark$	$[\text{OH}^-] = 0,01 \text{ mol}\cdot\text{dm}^{-3} \checkmark$
$[\text{OH}^-] = 0,01 \text{ mol}\cdot\text{dm}^{-3} \checkmark$	

(4)

7.2.3 Positive marking from / Positiewe nasien vanaf 7.2.1 and/ en 7.2.2  
Marking criteria / Nasienkriteria

- Using ratio Acid : Base 1 : 1
- Subst. of base values into  $n_{\text{NaOH excess}} = cV$
- Addition of n remaining and n initial (NaOH)
- Subst. into  $c = n/V$
- Multiplication of  $c_{\text{dilute}}$  by 10
- Final answer
- **Gebruik verhouding Suur : Basis = 1 : 1**
- **Vervanging van basis waarde in  $n_{\text{NaOH oormaat}} = cV$**
- **Addisie van n oorbly en n aanvanklik (NaOH)**
- **Vervanging in  $c = n / V$**
- **Vermenigvuldiging van  $c_{\text{verdun}}$  met 10**
- **Finale antwoord**

$$n(\text{NaOH})_{\text{reacting/reageer}} = 2,5 \times 10^{-3} \text{ mol} \checkmark \text{ From/Vanaf 7.2.1}$$

$$n(\text{NaOH})_{\text{in excess/oormaat}} = cV$$

$$n(\text{NaOH})_{\text{excess/oormaat}} = (0,01)(60 \times 10^{-3}) \checkmark$$

$$n(\text{NaOH})_{\text{excess/oormaat}} = 6 \times 10^{-4} \text{ mol}$$

$$n(\text{NaOH})_{\text{total/totaal}} = 2,5 \times 10^{-3} + 6 \times 10^{-4} \checkmark$$

$$n(\text{NaOH}) = 3,1 \times 10^{-3} \text{ mol}$$

(6)

$$C_{\text{dilute/verdun}} = \frac{n}{V}$$

$$C_{\text{dilute/verdun}} = \frac{3,1 \times 10^{-3}}{35 \times 10^{-3}} \checkmark$$

$$C_{\text{dilute/verdun}} = 0,08857 \text{ mol} \cdot \text{dm}^{-3}$$

$$\begin{aligned} C_{\text{concentrated /}} &= 0,08857 \times 10 \checkmark \\ &= 0,8857 \text{ mol} \cdot \text{dm}^{-3} \checkmark / 0,89 \text{ mol} \cdot \text{dm}^{-3} \end{aligned}$$

[19]

**QUESTION 8/VRAAG 8**

8.1 Chemical energy is converted into electrical energy ✓✓  
Chemiese energie word na elektriese energie omgeskakel. (2)

8.2.1  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+} + \text{e}^-$  ✓✓ (2)

**Marking criteria / Nasienkriteria**

- $\text{Fe}^{3+} + \text{e}^- \leftarrow \text{Fe}^{2+}$  2/2
- $\text{Fe}^{2+} \rightleftharpoons \text{Fe}^{3+} + \text{e}^-$  ½
- $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$  0/2
- $\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$  0/2
  
- Ignore if the charge omitted on electron  
*Ignoreer indien lading op elektron weggelaat is*
  
- If a charge of an ion is omitted  
*As lading weggelaat is op 'n foon.*  
e.g. / bv.  $\text{Fe}^2 \rightarrow \text{Fe}^3 + \text{e}^-$  Max. / Maks. ½

8.2.2  $\text{Ag} | \text{Ag}^+ \vee | | \checkmark \text{Fe}^{3+}, \text{Fe}^{2+} \vee | \text{Pt} \vee$  (3)

**Marking criteria/Nasienkriteria**

- $\text{Ag} | \text{Ag}^+ \vee$
- $\text{Fe}^{3+}, \text{Fe}^{2+} | \text{Pt} \vee$
- || ✓

8.2.3 Concentration:  $1 \text{ mol}\cdot\text{dm}^{-3}$  ✓ and temperature:  $25^\circ\text{C}$  ✓ /  $298 \text{ K}$   
Konsentrasie:  $1 \text{ mol}\cdot\text{dm}^{-3}$  en temperatuur:  $25^\circ\text{C}$  /  $298 \text{ K}$  (2)

8.3  $E^\theta_{\text{cell}} = E^\theta_{\text{cathode/reduction/oxidising agent}} - E^\theta_{\text{anode/oxidation/reducing agent}}$  ✓

$$E^\theta_{\text{cell}} = (0,80) \vee - (0,77) \vee$$

$$E^\theta_{\text{cell}} = 0,03 \text{ V} \vee$$

**Marking criteria/Nasienkriteria**

- Any other formula using unconventional abbreviation , e.g.  
*Enige ander formule wat onkonvensionele afkortings gebruik bv.*
- $E^\theta_{\text{cell}} = E^\theta_{\text{OA}} - E^\theta_{\text{RA}}$  followed by the correct substitution./ gevvolg deur korrekte vervangings ¾

(4)

8.4 Decrease / Afneem ✓ (1)

8.5  $\text{Cl}^-$  would form an insoluble salt/precipitate with the  $\text{Ag}^+$  ✓✓ /  $\text{AgCl}$  will precipitate out. The  $\text{Ag}^+$  half-cell would no longer be neutral / Circuit would be incomplete / Not enough electrolyte in half cell.

$\text{Cl}^-$  sal 'n neerslag vorm/presipiteer met  $\text{Ag}^+$ /  $\text{AgCl}$  vorm 'n neerslag / Die  $\text{Ag}^+$ halfsel sal nie meer neutraal wees nie / Stroombaan sal onvoltooied wees / Nie genoeg elektroliet in die halfsel nie. (2)

[16]

## QUESTION 9/VRAAG 9

9.1

**Marking criteria/ Nasienriglyne**

If any of the underlined key words/phrases in the **correct context** are omitted: - 1 mark per word/phrase.

*Indien enige van die sleutelwoorde/frases in die korrekte konteks weggelaat word: - 1 punt per woord/frase*

The chemical process in which electrical energy is converted to chemical energy. ✓✓

*Die chemiese proses waarin elektriese energie word na chemiese energie omgeskakel.*

**OR / OF**

The use of electrical energy to produce a chemical change. ✓✓

*Die gebruik van elektriese energie om chemiese energie te produseer* (2)

9.2

ENDOTHERMIC / ENDOTERMIES ✓

(1)

9.3

 $2 \text{Cl}^- \rightarrow \text{Cl}_2(\text{g}) + 2\text{e}^-$  ✓✓

Ignore phases / Ignoreer fases

(2)

**Marking criteria / Nasienkriteria**

- $2 \text{Cl}^- \rightleftharpoons \text{Cl}_2(\text{g}) + 2\text{e}^- \quad \frac{1}{2}$
- $\text{Cl}_2(\text{g}) + 2\text{e}^- \leftarrow 2 \text{Cl}^- \quad 2/2$
- $\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2 \text{Cl}^- \quad 0/2$

Ignore if the charge omitted on electron /

*Ignoreer as lading op elektron weggelaat is*

9.4

$$n = \frac{m}{M}$$

$$n = \frac{0,369}{63,5} \checkmark$$

$$n = 5,811 \times 10^{-3}$$

$$n (e^-) = 2 \times 5,811 \times 10^{-3} \checkmark$$

$$n (e^-) = 0,01162$$

$$n = \frac{N}{N_A}$$

$$0,01162 = \frac{N}{6,02 \times 10^{23}} \checkmark$$

$$N = 6,9964 \times 10^{21}$$

$$Q = Nq$$

$$Q = 6,9964 \times 10^{21} \times 1,6 \times 10^{-19} \checkmark$$

$$Q = 1119,4310 \text{ C}$$

$$I = \frac{Q}{\Delta t} \checkmark$$

$$I = \frac{1119,4310}{27 \times 60} \checkmark$$

$$I = 0,69 \text{ A} \checkmark$$

(7)  
[12]**TOTAL/TOTAAL:** 150**Marking criteria/Nasienkriteria**

- Subst. into  $n = m/M$
- Use of mole ratio Cu :  $e^-$
- Subst. into  $n = N/N_A$
- Subst. into  $n = Q/q_e$
- Formula  $Q = I\Delta t$
- Subst. into  $Q = I\Delta t$
- Final answer with unit

- *Vervang in n = m/M*
- *Gebruik van mol-verhouding Cu :  $e^-$*
- *Vervang in n = N/N\_A*
- *Vervang in n = Q/q\_e*
- *Formule Q = IΔt*
- *Vervang in Q = IΔt*
- *Finale antwoord met eenheid*